



Eurythmy Therapy

Focus on Movement for Better Health

A LOOK INTO THE ANTHROPOSOPHIC MEDICAL SYSTEM



TABLE OF CONTENTS

Eurythmy Therapy – what is it.....	pg 01
Qualifications and settings.....	pg 01
Mindfulness as a curative impact in Eurythmy therapy.....	pg 02
Reported effects of Eurythmy therapy.....	pg 03
The individual case in research – the Case Vignette Eurythmy Therapy (CAVI-YET).....	pg 04
References.....	pg 05

Eurythmy Therapy – what it is



Eurythmy therapy (EuTh) is a holistic, **self-activating mind-body** intervention within Anthroposophic Medicine that uses movement exercises with the arms, legs and the whole body. The exercises convert the **sounds of speech and music into movement**. The purpose of EuTh is to promote self-regulation and activate the **self-healing forces** in the human organism. Its integrative medical approach is based on conventional medicine, and in addition to the physical constitution of the patients, it also focuses on their **soul, spirit and life processes**. Eurythmy therapy exercises are recommended for specific symptoms, pathologies and constitutional peculiarities.

Qualifications and settings

Following a 4-year basic Eurythmy training, Eurythmy therapists undergo a full- or part-time therapy training¹, which includes general medical knowledge; training on the anthroposophic understanding of man and nature, health and illness; specific competences for the therapeutic discipline, with mentored practice and project work and medical-therapeutic research. Eurythmy therapists receive a diploma or a Masters' degree after presenting a case report and completing a 6-month internship after the training. Continued professional development is required to maintain a professional level.

Eurythmy therapy is widely implemented in anthroposophic outpatient and clinical settings, both in group or individual therapy. It is also practiced in pre-schools and schools. Eurythmy therapy is generally prescribed by a health care professional, but can also be recommended by a teacher or sought out by the patients themselves. The therapy plan is based on the medical diagnosis, and the movement diagnosis by the therapist. Over the course of 6-12 sessions, the patient learns how to perform the exercises, which she/he has to apply at home: regular independent practice reinforces the therapeutic effect [1], as the patient learns to care for him/herself [2].




¹ 7.200 h basic Eurythmy and 1.800 h Eurythmy therapy training



Positive impact of mindfulness aspects in Eurythmy therapy

Eurythmy therapy combines mindfulness components with imaginative movement components. Through movement, the external image becomes an internally experienced, healthy pre-image [7, pp. 128–129]. This can lead to a slowing down of movement [3] and to a heart rate variability improvement (HRV)[4]. Slowing down is usually associated with greater mindfulness, which in turn has a positive influence on psychological factors, such as in patients with inflammatory bowel disease [5] [6].[8] [9] [10].

As displayed in a questionnaire by Arndt Büssing, a correlation between the reduction in symptoms and mental stabilization is observed while practicing EuTh [11].



Reported effects of Eurythmy therapy

An overview of the effectiveness of Eurythmy therapy in clinical settings from 1985 onwards has been provided by Arndt Büssing et al. [12]. In these studies, EuTh as part of the treatment of **heart attack, attention deficit hyperactivity disorder (ADHD), anorexia nervosa, depression, anxiety, chronic diseases, such as asthma, migraine, low back pain**, and others was reported to show significant improvement in symptoms in clinically relevant trial sizes. Lötzke et al. presented a second systematic literature review, which confirmed the above results and noted positive treatment effects also for other conditions: **essential hypertension, posterior fossa tumor, status before and after surgery, moderate stress, fatigue and anxiety** [13].

GLOBALLY



523.2 million

people are living with
cardiovascular disease

5 %

of children and
adolescence have ADHD

280 million

people suffer from
depression

**Statistic sources are listed at the end of the 'References' page (page 6)*



Since then, more research has been done. An important treatment field is oncology, where **cancer related fatigue (CRF)** is well researched [14][15][16] [17]. Further impact, such as **mitigating side effects of chemotherapy and dealing with anxiety and companionship in the palliative situation** are highlighted by Barbara Trapp [18]. A new comparative RCT (EuTh vs. slow movement fitness) is currently running on CRF [19]. **Heart Rate Variability (HRV) improvement** through Eurythmy therapy can be observed from several of the studies mentioned above [8] [9] [10]. The effect of Eurythmy therapy on **fatigue** [20], **stress** [21] [22], **depression** [23] [24] and **anxiety** [25] [26] is described not only in terms of symptom reduction, but also on increasing health-related quality of life (HRQL) indexes. That effect was also described in a comparative RCT on **chronic back pain** in the YES study [27]: reduction of pain was reported from EuTh, as well as from yoga and physiotherapy; HRQL was clearly stronger in the EuTh-participants in a 16-week follow up. Regarding the effect on children, a routine outcome monitoring

done in the Netherlands shows statistically significant improvements from eurythmy therapy-specific observation items in children aged 3-18 [28]. Annette Weißkircher presented excellent results in the prevention and treatment of hay fever [29].

The individual case in research – the Case Vignette Eurythmy Therapy (CAVI-EYT)

Because of the holistic and patient-centered approach, CAVI-EYT can give a comprehensive view of the treatment process with all the facets that are typically lost in RCTs, and can provide indications of the effectiveness of the therapy for treatment practice. In order to publish more cases, a CAVI-EYT guideline was developed in 2020 [30][31][32]. It is a structured presentation of individual case treatments with eurythmy therapy in an abridged version. The summary is based on the CARE Guideline [33] and is intended to enable therapists without detailed medical knowledge to present their cases appropriately and in a way that is relevant to the scientific discourse. CAVI-EYT was presented to international committees and has since been used to publish EuTh case vignettes [34] [23] [24] [35] [2].



REFERENCES

- [1]K. Gerlach and U. Pohlman, 'Abklingen von Migräne-Symptomatik nach Eurythmietherapie', *Merkurstab*, vol. 70, no. 4, pp. 312–317, 2017, doi: 10.14271/DMS-20821-DE.
- [2]K. Gerlach, 'Patient Reported Effects of self-help Eurythmy Therapy managing Covid-19-Illness "A Case-Vignette', *J. Complement. Med. Res.*, vol. 13, no. 1, p. 122, 2022, doi: 10.5455/jcmr.2022.13.01.22.
- [3]F. Edelhäuser, M. Heidelbach, B. Trapp, and D. Cysarz, 'Wirkmechanismen der Eurythmietherapie – Ergebnisse aktueller Forschung', *Merkurstab*, vol. 74, no. 1, pp. 4–16, 2021, doi: 10.14271/DMS-21304-DE.
- [4]F. Edelhäuser, A. Minnerop, B. Trapp, A. Büssing, and D. Cysarz, 'Eurythmy therapy increases specific oscillations of heart rate variability', *BMC Complement. Altern. Med.*, vol. 15, no. 1, Dec. 2015, doi: 10.1186/s12906-015-0684-6.
- [5]M. M. Hood and S. Jedel, 'Mindfulness-Based Interventions in Inflammatory Bowel Disease', *Gastroenterol. Clin. North Am.*, vol. 46, no. 4, pp. 859–874, Dec. 2017, doi: 10.1016/j.gtc.2017.08.008.
- [6]T. Ewais et al., 'A systematic review and meta-analysis of mindfulness based interventions and yoga in inflammatory bowel disease', *J. Psychosom. Res.*, vol. 116, pp. 44–53, Jan. 2019, doi: 10.1016/j.jpsychores.2018.11.010.
- [7]R. Steiner, *Geisteswissenschaftliche Gesichtspunkte zur Therapie. Heileurythmie Zwei Kurse in chronologischer Folge*. Rudolf Steiner Verlag, 2016.
- [8]G. Seifert et al., 'Effects of complementary eurythmy therapy on heart rate variability', *Complement. Ther. Med.*, vol. 17, no. 3, pp. 161–167, Jun. 2009, doi: 10.1016/j.ctim.2008.09.005.
- [9]G. Seifert et al., 'Improvement of Heart Rate Variability by Eurythmy Therapy After a 6-Week Eurythmy Therapy Training', *Integr. Cancer Ther.*, vol. 11, no. 2, pp. 111–119, Jun. 2012, doi: 10.1177/1534735411413263.
- [10]G. Seifert et al., 'Improvement of Circadian Rhythm of Heart Rate Variability by Eurythmy Therapy Training', *Evid. Based Complement. Alternat. Med.*, vol. 2013, pp. 1–9, 2013, doi: 10.1155/2013/564340.
- [11]A. Büssing, F. Edelhäuser, A. Weisskircher, J. M. Fouladbakhsh, and P. Heusser, 'Inner Correspondence and Peacefulness with Practices among Participants in Eurythmy Therapy and Yoga: A Validation Study', *Evid. Based Complement. Alternat. Med.*, vol. 2011, pp. 1–9, 2011, doi: 10.1155/2011/329023.
- [12]A. Büssing, T. Ostermann, M. Majorek, and P. F. Matthiessen, 'Eurythmy Therapy in clinical studies: a systematic literature review', *BMC Complement. Altern. Med.*, vol. 8, no. 1, Dec. 2008, doi: 10.1186/1472-6882-8-8.
- [13]D. Lötze, P. Heusser, and A. Büssing, 'A systematic literature review on the effectiveness of eurythmy therapy', *J. Integr. Med.*, vol. 13, no. 4, pp. 217–230, Jul. 2015, doi: 10.1016/S2095-4964(15)60163-7.
- [14]G. S. Kienle, H. Kiene, and H. U. Albonico, 'Anthroposophische Medizin: Health Technology Assessment Bericht – Kurzfassung', *Forsch. Komplementärmedizin Res. Complement. Med.*, vol. 13, no. 2, pp. 7–18, 2006, doi: 10.1159/000093481.
- [15]G. S. Kienle, H.-U. Albonico, E. Baars, H. J. Hamre, P. Zimmermann, and H. Kiene, 'Anthroposophic Medicine: An Integrative Medical System Originating in Europe', *Glob. Adv. Health Med.*, vol. 2, no. 6, pp. 20–31, Nov. 2013, doi: 10.7453/gahmj.2012.087.
- [16]M. Kröz et al., 'Multimodal Therapy Concept and Aerobic Training in Breast Cancer Patients With Chronic Cancer-Related Fatigue', *Integr. Cancer Ther.*, vol. 12, no. 4, pp. 301–311, Jul. 2013, doi: 10.1177/1534735412464552.
- [17]G. Bräuner-Gülow, 'Dokumentation zur Heileurythmie bei Mamma-Karzinom', *Merkurstab*, 2008, doi: 10.14271/DMS-19224-DE.
- [18]B. Trapp, 'Heileurythmie als umfassende Begleitung für Patienten mit einer malignen Tumorerkrankung', *Merkurstab*, vol. 70, no. 3, pp. 237–240, 2017, doi: 10.14271/DMS-20797-DE.
- [19]D. Meier-Girard, K. Ribbi, G. Gerstenberg, T. Ruhstaller, and U. Wolf, 'Eurythmy therapy versus slow movement fitness in the treatment of fatigue in metastatic breast cancer patients: study protocol for a randomized controlled trial', *Trials*, vol. 21, no. 1, p. 612, Dec. 2020, doi: 10.1186/s13063-020-04542-5.
- [20]J. L. Kanitz et al., 'The impact of eurythmy therapy on fatigue in healthy adults—A controlled trial', *Eur. J. Integr. Med.*, vol. 4, no. 3, pp. e289–e297, Sep. 2012, doi: 10.1016/j.eujim.2012.02.006.
- [21]J. L. Kanitz et al., 'The impact of eurythmy therapy on stress coping strategies and health-related quality of life in healthy, moderately stressed adults', *Complement. Ther. Med.*, vol. 19, no. 5, pp. 247–255, Oct. 2011, doi: 10.1016/j.ctim.2011.06.008.
- [22]B. Berger, M. Bertram, J. Kanitz, K. Pretzer, and G. Seifert, "'Like Walking into an Empty Room": Effects of Eurythmy Therapy on Stress Perception in Comparison with a Sports Intervention from the Subjects' Perspective—A Qualitative Study', *Evid. Based Complement. Alternat. Med.*, vol. 2015, pp. 1–11, 2015, doi: 10.1155/2015/856107.
- [23]K. Gerlach, V. Veit, S. Walther, and A. Weißkircher, 'Behandlung einer milden Depression und eines psychosomatischen

Reizdarmsyndroms mit achtsamkeitsorientierter Eurythmietherapie – Eine Fallvignette’, *Merkurstab*, vol. 74, no. 4, pp. 355–359, 2021, doi: 10.14271/DMS-21386-DE.

[24]I. Hermansen and D. McGavin, ‘Reduction of psychotropic and analgesic medication using Eurythmy Therapeutic vowel A for a patient with multimorbidity – a case vignette’, *Merkurstab*, vol. 74, no. 5, pp. 417–420, 2021, doi: 10.14271/DMS-21408-EN.

[25]J. Schwab et al., ‘Eurythmy Therapy in anxiety’, *Altern. Ther. Health Med.*, vol. 17, no. 4, pp. 56–63, Aug. 2011.

[26]K. Gerlach, ‘Prozessorientierung in der Heileurythmie – Kasuistik zu einer Behandlung von Fahrangst’, *Merkurstab*, vol. 74, no. 1, pp. 54–62, 2021, doi: 10.14271/DMS-21309-DE.

[27]A. Büssing, D. Poier, T. Ostermann, M. Kröz, and A. Michalsen, ‘Treatment of Chronic Lower Back Pain: Study Protocol of a Comparative Effectiveness Study on Yoga, Eurythmy Therapy, and Physiotherapeutic Exercises’, *Complement. Med. Res.*, Jul. 2017, doi: 10.1159/000471801.

[28]E. Baars, N. van Steenberghe, R. de Zwaan, A. Kraakman, M. de Vries, and M. Wodowoz- de Boon, ‘Effecten van en ervaringen met Eurythmietherapie voor kinderen’, Leiden, 2021. [Online]. Available: https://www.hsleiden.nl/binaries/content/assets/hsl/lectoraten/antroposofische-gezondheidszorg/publicaties/boeken_rapporten/effecten-van-en-ervaringen-met-eurythmie-voor-kinderen

[29]P. Ehrler et al., ‘Die Wirksamkeit der heileurythmischen Therapielautreihe TSRMA bei allergischer Rhinopathie – Eine qualitative Inhaltsanalyse nach Mayring’, *Merkurstab*, vol. 74, no. 1, pp. 37–44, 2021, doi: 10.14271/DMS-21307-DE.

[30]K. Gerlach, ‘The Case-Vignette in Eurythmy Therapie CA VI-EYT’, Jan. 2021. https://eurythmytherapy-medsektion.net/fileadmin/user_upload/pdf/FAVI_6-2-2021_pptx_compressed.pdf

[31]K. Gerlach, ‘Leitfaden für Fallvignetten in der Eurythmietherapie’, *Merkurstab*, vol. 74, no. 3, p. 284, 2021, doi: 10.14271/DMS-21367-DE.

[32]‘Abstracts der Poster der Jahreskonferenz der Medizinischen Sektion am Goetheanum, 17.–19.9.2021, Dornach (Teil 1)’, *Merkurstab*, vol. 75, no. 1, pp. 56–61, 2022, doi: 10.14271/DMS-21458-DE.

[33]J. J. Gagnier, G. Kienle, D. G. Altman, D. Moher, H. Sox, and D. Riley, ‘The CARE Guidelines: Consensus-based Clinical Case Reporting Guideline Development’, *Glob. Adv. Health Med.*, vol. 2, no. 5, pp. 38–43, Sep. 2013, doi: 10.7453/gahmj.2013.008.

[34]K. Gerlach, ‘Belastung durch Post-Zoster-Neuralgie mit Reizblase und Augenlidzucken behandelt mit Eurythmietherapie – Eine Fallvignette’, *Merkurstab*, vol. 74, no. 2, pp. 163–166, 2021, doi: 10.14271/DMS-21336-DE.

[35]I. Stadler, K. Gerlach, V. Veit, and M. Krüger, ‘Behandlung eines Morbus Crohn mit Eurythmietherapie – Eine Fallvignette’, *Merkurstab*, vol. 74, no. 6, pp. 490–495, 2021, doi: 10.14271/DMS-21427-DE.

Photos source:

All photos used in the paper are taken by Till Kleinelützum for Katharina Gerlach.

Statistics source (pg.3):

1. American Heart Association Report ‘*Heart disease #1 cause of death rank likely to be impacted by COVID-19 for years to come*’; January 2021; available at <https://newsroom.heart.org/news/heart-disease>

2. National Library of Medicine, ‘*The prevalence of adult attention-deficit hyperactivity disorder: A global systematic review and meta-analysis*’; February 2021; available at: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7916320/#:~:text=Globally%2C%20it%20has%20been%20estimated,affected%20by%20ADHD%20%5B10%5D>.

3. World Health Organization: Depression Factsheet from September 2021; available at <https://www.who.int/news-room/factsheets/detail/depression>

This paper is part of a series ‘A Look into the Anthroposophic Medical System’, which explores the different professions within the anthroposophic medicine field. For more information, please contact hana.adamcova@medsektion-ikam.org or contact@ivaa.info.



100 years
Eurythmy Therapy

International coordination, Eurythmy Therapy
www.eurythmytherapy-medsektion.net



International Federation
of Anthroposophic
Medical Associations

www.ivaa.info